

Non-perturbative $O(a)$ -improvement of Wilson quark action in three-flavor QCD with plaquette gauge action

N. Yamada¹, S. Aoki², M. Fukugita³, S. Hashimoto¹, K-I. Ishikawa⁴,
N. Ishizuka^{2,5}, Y. Iwasaki^{2,5}, K. Kanaya^{2,5}, T. Kaneko¹, Y. Kuramashi^{2,5},
M. Okawa⁴, Y. Taniguchi², N. Tsutsui¹, A. Ukawa^{2,5}, and T. Yoshié^{2,5}
(CP-PACS and JLQCD Collaborations)

¹*High Energy Accelerator Research Organization(KEK), Tsukuba, Ibaraki 305-0801, Japan*

²*Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan*

³*Institute for Cosmic Ray Research, University of Tokyo, Kashiwa 277-8582, Japan*

⁴*Department of Physics, Hiroshima University,
Higashi-Hiroshima, Hiroshima 739-8526, Japan*

⁵*Center for Computational Sciences, University of Tsukuba, Tsukuba, Ibaraki 305-8577, Japan*

(Dated: February 1, 2008)

We perform a non-perturbative determination of the $O(a)$ -improvement coefficient c_{SW} for the Wilson quark action in three-flavor QCD with the plaquette gauge action. Numerical simulations are carried out in a range of $\beta=12.0-5.2$ on a single lattice size of $8^3 \times 16$ employing the Schrödinger functional setup of lattice QCD. As our main result, we obtain an interpolation formula for c_{SW} and the critical hopping parameter K_c as a function of the bare coupling. This enables us to remove $O(a)$ scaling violation from physical observables in future numerical simulation in the wide range of β . Our analysis with a perturbatively modified improvement condition for c_{SW} suggests that finite volume effects in c_{SW} are not large on the $8^3 \times 16$ lattice. We investigate N_f dependence of c_{SW} by additional simulations for $N_f = 4, 2$ and 0 at $\beta=9.6$. As a preparatory step for this study, we also determine c_{SW} in two-flavor QCD at $\beta=5.2$. At this β , several groups carried out large-scale calculations of the hadron spectrum, while no systematic determination of c_{SW} has been performed.

I. INTRODUCTION

Recent large-scale simulations in two-flavor QCD have demonstrated that quenching artifacts found in various physical observables are reduced by dynamical effects of up and down quarks. There has been significant progress also in the algorithms for QCD with odd numbers of flavors: while the conventional Hybrid Monte Carlo (HMC) algorithm [1] is applicable only to even-flavor QCD, the exact algorithms, such as the Multi-Boson [2] and the Polynomial HMC algorithms [3] capable of odd-flavor cases, have been developed. Clearly the time has come to undertake a fully realistic and extensive simulations of QCD with dynamical up, down and strange quarks.

Since simulations with dynamical quarks are computationally demanding, highly improved lattice actions should be employed in the three-flavor simulations. The leading cutoff effect in physical quantities is $O(a)$ with the Wilson quark action, and this error can be removed by adding a single counter term, the Sheikholeslami-Wohlert (SW) term [4], to the action with non-perturbatively determined coefficient c_{SW} . However, c_{SW} has been determined only in quenched and two-flavor QCD so far [5, 6].

In this article, we perform a non-perturbative determination of c_{SW} in three-flavor QCD with the plaquette gauge and the Wilson quark actions. In Refs. [7, 8], however, we found that this theory has a non-trivial phase structure: there is an unphysical phase transition at $\beta \approx 5.0$, where the lattice cutoff a^{-1} is roughly 2.6 GeV. It is expected that the phase transition strongly distorts scaling properties of physical observables. This transition is considered as an artifact due to the finite lattice spacing and can be removed by the use of improved gauge actions [7, 8]. Therefore, there are two strategies for meaningful simulations in three-flavor QCD: i) use the plaquette gauge action at $\beta \gg 5.0$, or ii) use an improved gauge action, if simulations at $a^{-1} \lesssim 2.6$ GeV are needed. We explore the former strategy in this article, and the latter possibility is studied in a separate publication [9].

In our determination of c_{SW} , we follow the method proposed by the ALPHA Collaboration in Refs. [5, 10]. We explore a wide range of $\beta=12.0-5.2$, which is significantly higher than the phase transition point $\beta \approx 5.0$, employing a single lattice size of $8^3 \times 16$. As our main result, we derive an interpolation formula for c_{SW} as a function of the bare coupling. The critical hopping parameter K_c in the $O(a)$ -improved theory is also obtained

as a byproduct. We examine finite volume effects in c_{SW} by modifying the improvement condition at one-loop order of perturbation theory. Additional simulations in four-, two-flavor and quenched QCD at $\beta = 9.6$ are carried out to investigate the N_f dependence of c_{SW} .

As a preparatory step toward this study, we also determine c_{SW} in two-flavor QCD at $\beta = 5.2$. In the previous work by the ALPHA Collaboration [6], they carried out the non-perturbative tuning of c_{SW} at $\beta \geq 5.4$, and derived an interpolation formula of their result as a function of g_0^2 . However, due to the limitation of available computer power, recent large-scale simulations by the UKQCD[11] and JLQCD Collaborations[12] were performed at a lower value $\beta = 5.2$ with c_{SW} obtained by extrapolating the ALPHA's formula. We, therefore, determine c_{SW} directly at $\beta = 5.2$ in order to see if the extrapolation of the formula to this value of β really works out, and to confirm that $O(a)$ scaling violation is absent in the UKQCD and JLQCD simulations.

This paper is organized as follows. In Sec.II, we briefly introduce the method for the non-perturbative tuning of c_{SW} employed in this study. Section III is devoted to detailed description of our numerical analysis and results in two-flavor QCD at $\beta = 5.2$. We present our results in three-flavor QCD and discuss its $O(a/L)$ uncertainty in Sec. IV. Finally our conclusion is given in Sec. V.

II. IMPROVEMENT CONDITION FOR $O(a)$ -IMPROVEMENT

In our determination of c_{SW} , we basically follow the method proposed in Refs. [5, 10], which employs the Schrödinger functional (SF) setup of lattice QCD [13]. In this section, we briefly introduce the SF setup and the choice of the improvement condition to fix c_{SW} .

A. SF setup

The SF is the generating functional of the field theory with the Dirichlet boundary condition imposed in the temporal direction. In this study, the spatial link variables at the boundaries are set to the following diagonal, constant $SU(3)$ matrices

$$U_k(\mathbf{x}, x_0)|_{x_0=0} = \exp[aC_k], \quad U_k(\mathbf{x}, x_0)|_{x_0=T} = \exp[aC'_k], \quad (1)$$

$$C_k = \frac{i\pi}{6L_k} \begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad C'_k = \frac{i\pi}{6L_k} \begin{pmatrix} -5 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad (2)$$

where L_k ($k=1, 2, 3$) and T are physical lattice sizes in the spatial and temporal directions. All quark fields at the boundaries are set to zero. In the spatial directions, the periodic boundary condition is imposed for both gauge and quark fields.

We use the plaquette gauge action

$$S_g = \frac{\beta}{6} \sum_{x,\mu,\nu} \text{Tr} [1 - U_{x,\mu\nu}], \quad (3)$$

where $U_{x,\mu\nu}$ is the product of gauge link variables $U_{x,\mu}$ around the plaquette

$$U_{x,\mu\nu} = U_{x,\mu} U_{x+\hat{\mu},\nu} U_{x+\hat{\nu},\mu}^\dagger U_{x,\nu}^\dagger. \quad (4)$$

The $O(a)$ -improved Wilson quark action [4] is given by

$$S_q = \sum_{x,y} \bar{q}_x D_{xy} q_y, \quad (5)$$

$$D_{xy} = \delta_{xy} - K \sum_{\mu} \left\{ (1 - \gamma_{\mu}) U_{x,\mu} \delta_{x+\hat{\mu},y} + (1 + \gamma_{\mu}) U_{x-\hat{\mu},\mu}^\dagger \delta_{x-\hat{\mu},y} \right\} + \frac{i}{2} K c_{\text{SW}} \sigma_{\mu\nu} F_{x,\mu\nu} \delta_{xy}, \quad (6)$$

with the field strength tensor $F_{x,\mu\nu}$ defined by

$$F_{x,\mu\nu} = \frac{1}{8} \{ (U_{x,\mu\nu} + U_{x,\nu-\mu} + U_{x,-\mu-\nu} + U_{x,-\nu\mu}) - (\text{h.c.}) \}, \quad (7)$$

where (h.c.) denotes the hermitian conjugate of the preceding bracket, and $\sigma_{\mu\nu} = (i/2) [\gamma_\mu, \gamma_\nu]$. The last term in Eq. (6) is the counter term to remove $O(a)$ effects in on-shell quantities. Its coefficient c_{SW} is set to unity to remove tree-level $O(a)$ scaling violation from physical observables. The main purpose in this article is non-perturbative tuning of c_{SW} for removal of all $O(ag_0^n)$ scaling violation ($n \geq 0$). For the $O(a)$ -improvement of the SF itself, we add counter terms made of the gauge and quark fields at boundaries to the lattice action. However, these counter terms affect the PCAC relation at order of a^2 or higher, and hence are not necessary for determination of c_{SW} from the PCAC relation. In the ALPHA Collaboration's studies, the counter terms are omitted except for a term

$$\delta S_g = \frac{\beta}{6} (c_t - 1) \sum_{\mathbf{x}, \mu} \text{Tr} [1 - U_{x, \mu 0}], \quad (8)$$

which is made of the temporal plaquettes touching the boundaries. In this study, we also include this counter term to the total lattice action $S = S_g + \delta S_g + S_q$ so that we can directly compare our and ALPHA Collaboration's results. The coefficient of the counter term c_t is set to the one-loop estimate in Ref. [14].

B. Improvement condition

We determine c_{SW} by imposing the validity of the PCAC relation

$$\frac{1}{2} (\partial_\mu + \partial_\mu^*) A_{\text{imp}, \mu}^a = 2mP^a, \quad (9)$$

up to order of a^2 . The pseudo-scalar operator and $O(a)$ -improved and unimproved axial currents are given by

$$P^a = \bar{\psi} \gamma_5 \tau^a \psi, \quad (10)$$

$$A_{\text{imp}, \mu}^a = A_\mu^a + c_A \frac{1}{2} (\partial_\mu + \partial_\mu^*) P^a, \quad (11)$$

$$A_\mu^a = \bar{\psi} \gamma_\mu \gamma_5 \tau^a \psi, \quad (12)$$

where ∂_μ and ∂_μ^* are the forward and backward lattice derivatives and $SU(N_f)$ generators τ^a act on the flavor indices of the quark fields $\bar{\psi}$ and ψ .

We measure two correlation functions

$$f_A(x_0) = -\frac{1}{N_f^2 - 1} \langle A_0^a(x) \mathcal{O}^a \rangle, \quad (13)$$

$$f_P(x_0) = -\frac{1}{N_f^2 - 1} \langle P^a(x) \mathcal{O}^a \rangle, \quad (14)$$

where $\langle \dots \rangle$ denotes the expectation value after taking trace over color and spinor indices and summing over spatial coordinate $\mathbf{x}$. For the source operator, we take

$$\mathcal{O}^a = a^6 \sum_{\mathbf{y}, \mathbf{z}} \bar{\zeta}(\mathbf{y}) \gamma_5 \tau^a \zeta(\mathbf{z}), \quad (15)$$

defined from the boundary fields

$$\zeta(\mathbf{x}) = \frac{\delta}{\delta \bar{\rho}(\mathbf{x})}, \quad \bar{\zeta}(\mathbf{x}) = \frac{\delta}{\delta \rho(\mathbf{x})}, \quad (16)$$

where $\rho(\mathbf{x})$ is the quark field at $x_0=0$ and is set to zero in the calculation of f_A and f_P . The bare quark mass is then calculated from f_A and f_P through the PCAC relation Eq. (9):

$$m(x_0) = r(x_0) + c_A s(x_0) \quad (17)$$

$$r(x_0) = \frac{1}{4} (\partial_0 + \partial_0^*) f_A(x_0) / f_P(x_0) \quad (18)$$

$$s(x_0) = \frac{1}{2} a \partial_0 \partial_0^* f_P(x_0) / f_P(x_0). \quad (19)$$

We can calculate another set of m' , r' and s' from the correlation functions

$$f'_A(T - x_0) = +\frac{1}{N_f^2 - 1} \langle A_0^a(x) \mathcal{O}'^a \rangle, \quad (20)$$

$$f'_P(T - x_0) = -\frac{1}{N_f^2 - 1} \langle P^a(x) \mathcal{O}'^a \rangle, \quad (21)$$

using the source operator at the other boundary

$$\mathcal{O}'^a = a^6 \sum_{\mathbf{y}, \mathbf{z}} \bar{\zeta}'(\mathbf{y}) \gamma_5 \tau^a \zeta'(\mathbf{z}), \quad (22)$$

where ζ' is the boundary field at $x_0 = T$.

The improvement condition to fix c_{SW} is obtained by requiring that quark masses calculated with different boundary conditions coincide with each other. However, a naive condition $m(x_0) = m'(x_0)$ requires a non-perturbative tuning of c_A as well as c_{SW} . To eliminate c_A from the process, it was proposed in Ref. [5] to use a modified definition of the quark mass

$$M(x_0, y_0) = m(x_0) - \frac{m(y_0) - m'(y_0)}{s(y_0) - s'(y_0)} s(x_0), \quad (23)$$

and similarly defined $M'(x_0, y_0)$. Therefore, c_{SW} is tuned so that the following mass difference

$$\Delta M(x_0, y_0) = M(x_0, y_0) - M'(x_0, y_0) \quad (24)$$

vanishes with a certain choice of (x_0, y_0) .

In principle, we can take an arbitrary choice for (x_0, y_0) , since a change of the choice leads to a difference in $O(a^2)$ scaling violation in physical observables. In this study, we take $(x_0, y_0) = (3T/4, T/4)$ for $\Delta M(x_0, y_0)$, and $(x_0, y_0) = (T/2, T/4)$ for $M(x_0, y_0)$. The latter is used to specify the massless point. We note that this choice is the same as that in the ALPHA's studies in quenched and two-flavor QCD. From now on, M and ΔM without arguments denote $M(T/2, T/4)$ and $\Delta M(3T/4, T/4)$, respectively.

In practice, c_{SW} is determined by demanding that M and ΔM satisfy the following improvement condition

$$\begin{cases} M &= 0, \\ \Delta M &= \Delta M^{(0)}, \end{cases} \quad (25)$$

where $\Delta M^{(0)}$ is the tree-level value of ΔM at the massless point $M=0$ on the finite lattice volume $L^3 \times T$. We tune ΔM to $\Delta M^{(0)}$ but not to zero so that the weak coupling limit of the non-perturbatively determined c_{SW} is exactly unity. On our lattice size of $8^3 \times 16$, $a\Delta M^{(0)} = 0.000277$ [5]. We also note that the tuning of M to the massless point provides a non-perturbative estimate of the critical hopping parameter K_c in the $O(a)$ -improved theory.

III. TWO-FLAVOR QCD AT $\beta=5.2$

A. Simulation method

In this section, we report the determination of c_{SW} in two-flavor QCD at $\beta=5.2$.

Our numerical simulations are carried out on a $8^3 \times 16$ lattice at six values of c_{SW} in a range $c_{\text{SW}} = 1.5 - 3.0$. We choose two to four values for the hopping parameter K at each c_{SW} so that we have data of ΔM at both of positive and negative values of M , and/or at M close to the massless point ($|aM| \leq 0.01$ in our study). This enables us to tune $(M, \Delta M)$ to $(0, \Delta M^{(0)})$ by an interpolation or short extrapolation. The simulated values of c_{SW} and K are summarized in Table I.

We use the standard HMC algorithm with the asymmetric even-odd preconditioning described in Refs. [15, 16, 17] for the determinant of the quark matrix D . We solve the linear equation $DX = B$ using the BiCGStab algorithm [18] with the stopping condition

$$\|R_i\|/\|B\| < 10^{-14}, \quad (26)$$

where $R_i = DX_i - B$ is the residual vector and X_i is the estimate for the solution X in the i -th BiCGStab iteration. The HMC trajectory length is fixed to the unit length. We set the number of the Molecular Dynamics steps to 60–80, which achieves the acceptance rate higher than 80%.

After the thermalization of 500 HMC trajectories, we accumulate the statistics N_{traj} summarized in Table I. The correlators f_X and f'_X ($X = A, P$) are measured at every trajectory. We use the jackknife method to estimate statistical errors of f_X , f'_X and all results derived from them.

B. numerical result

Numerical results of M and ΔM are summarized in Table II. In order to fix c_{SW} and K_c satisfying the improvement condition Eq. (25), we parameterize M and ΔM by a simultaneous fit in terms of $1/K$ and c_{SW} :

$$aM = a_M + \frac{b_M^{(1)}}{K} + \frac{b_M^{(2)}}{K^2} + c_M^{(1)} c_{\text{SW}} + c_M^{(2)} c_{\text{SW}}^2 + \frac{d_M}{K} c_{\text{SW}}, \quad (27)$$

$$a\Delta M = a_{\Delta M} + \frac{b_{\Delta M}^{(1)}}{K} + \frac{b_{\Delta M}^{(2)}}{K^2} + c_{\Delta M}^{(1)} c_{\text{SW}} + c_{\Delta M}^{(2)} c_{\text{SW}}^2 + \frac{d_{\Delta M}}{K} c_{\text{SW}}. \quad (28)$$

Fit parameters are summarized in Table III. Figure 1 shows $1/K$ dependence of M and ΔM , and M dependence of ΔM at each c_{SW} . By interpolating $(M, \Delta M)$ to $(0, \Delta M^{(0)})$ with this parameterization, we obtain

$$c_{\text{SW}} = 1.908(64), \quad (29)$$

$$K_c = 0.1381(12). \quad (30)$$

We also test another method for the parametrization of M and ΔM in order to estimate the systematic error due to the simultaneous fit Eqs. (27) and (28). At each c_{SW} , we determine ΔM at $M=0$ by a linear fit

$$a\Delta M = a'_M + b'_M(aM). \quad (31)$$

At $c_{\text{SW}}=1.90$ and 2.02 where we simulate more than two values for K , we also test a quadratic form and find that the higher order contribution is small and can be safely neglected in this analysis. Figure 2 shows c_{SW} dependence of ΔM at $M=0$ which we parametrize by linear or quadratic forms

$$a\Delta M = a'_{\Delta M} + b'_{\Delta M} c_{\text{SW}} + c'_{\Delta M} c_{\text{SW}}^2. \quad (32)$$

By tuning ΔM to its tree-level value, we obtain $c_{\text{SW}} = 1.979(68)$ from the linear fit as reported in Ref. [12], and $1.975(50)$ from the quadratic one. These are consistent with the result from the combined fit. This good agreement originates from our careful choice of c_{SW} and K in simulations: since we choose these parameters so that the region of $(M, \Delta M)$ contains $(0, \Delta M^{(0)})$ as shown in Fig. 1, c_{SW} can be fixed by a short interpolation for which the uncertainty due to the choice of the parameterization function for M and ΔM is not large.

The ALPHA's interpolation formula in Ref. [6] gives $c_{\text{SW}} = 2.017$ at $\beta = 5.2$, which is consistent with our results. This confirms that the ALPHA's formula can be used down to $\beta = 5.2$ as in the UKQCD[11] and JLQCD simulations[12].

However, as pointed out in Ref. [19], there are large cutoff effects in the PCAC quark mass m and the mass dependence of the Sommer scale r_0 [20] around $\beta = 5.2$. There is a possibility that the improvement condition Eq. (25) adopted in this and ALPHA's previous studies leaves unexpectedly large $O(a^2)$ scaling violations in physical observables around this value of β . Therefore, a test of alternative improvement conditions and scaling properties of physical observables is an important subject to avoid the large cutoff effects in future lattice calculations.

IV. THREE-FLAVOR QCD

A. Simulation method

We determine c_{SW} in three-flavor QCD at nine values of β in the range $\beta = 12.0 - 5.2$. Numerical simulations are carried out on a $8^3 \times 16$ lattice at four values of c_{SW} at each β , and three or four values of K at each c_{SW} .

These values are carefully chosen so that the region of $(M, \Delta M)$ contains or is sufficiently close to the point $(0, \Delta M^{(0)})$ which satisfies the improvement condition Eq. (25). These simulation parameters are summarized in Table IV.

In our simulations, we adopt the standard HMC algorithm for two-flavors of dynamical quarks and a polynomial HMC algorithm developed in Ref. [17] for the remaining one-flavor. We employ the symmetric even-odd preconditioning in Refs. [16, 17] for the quark matrix D . As in the two-flavor simulations at $\beta = 5.2$, we calculate D^{-1} using the BiCGStab algorithm with the tolerance parameter $\|R_i\|/\|B\| < 10^{-14}$. We set the number of the Molecular Dynamics steps to 80. This achieves the acceptance rate of about 90% or higher.

In the PHMC algorithm, we use the Chebyshev polynomial $P[D]$ to approximate D^{-1} . In order to make this algorithm exact, the correction factor

$$P_{\text{corr}} = \det[W[D]] \quad (33)$$

with $W[D] = P[D]D$ is taken into account by the noisy Metropolis method [21]. We calculate the square root of $W[D]$, which is required in the Metropolis test, with an accuracy of 10^{-14} using the Taylor expansion of $W[D]$ [17]. The order of the polynomial N_{poly} is chosen so that we achieve the acceptance rate of about 90 % or higher for the Metropolis test.

We note that, even with the SF setup, there is a difficulty in simulating massless or negative quark masses in three-flavor QCD. In the strong coupling region, eigenvalues of D have large fluctuations and they can take values outside the radius of convergence of $P[D]$. If this happens, the Polynomial approximation $P[D]$ and Taylor expansion of $W[D]$ break down. For this reason, our simulations in the strong coupling region are performed only down to $M \simeq 0$, while negative quark masses $M \simeq -0.03$ are simulated in the weak coupling region $\beta \simeq 12$.

We accumulate statistics N_{traj} summarized in Table IV, and measure the correlators f_X and f'_X ($X = A, P$) at every trajectory. The dependence of the jackknife error of M on the bin size N_{bin} is investigated in a range $N_{\text{bin}} = 1 - N_{\text{traj}}/20$. We then adopt N_{bin} giving the maximum error in the jackknife procedure in the following analysis.

We determine c_{sw} and K_c non-perturbatively also in four-flavor, two-flavor and quenched QCD at $\beta = 9.6$ to study their N_f dependence. The simulation method is similar to that in three-flavor QCD, except that we use the standard HMC algorithm in these cases. Simulation parameters are summarized in Tables V – VII.

B. Non-perturbative c_{sw} in three-flavor QCD

Numerical results of M and ΔM are summarized in Table VIII. In Fig. 3, we plot M and M' at several values of β as a function of x_0 . With our statistics, aM , aM' and hence $a\Delta M$ have an accuracy of 10^{-3} at all simulation parameters. These accurate data enables us to reduce the statistical error of c_{sw} to the level of $\sim 5\%$ even at our coarsest lattice spacing.

In order to parameterize the K and c_{sw} dependence of M and ΔM , we use the combined fit of Eqs. (27) and (28). Fit parameters are summarized in Tables IX and X. Figures 4 and 5 show M and ΔM and their fit curves at several values of β . We observe that our data are well described by the combined fit. Consequently, as shown in Fig. 6, M dependence of ΔM is reproduced reasonably well by the fit. We fix c_{sw} and K_c satisfying the improvement condition Eq. (25) using this parameterization. Numerical results for the non-perturbatively tuned c_{sw} and K_c are summarized in Table XI.

As in the analysis in two-flavor QCD at $\beta = 5.2$, we test Eqs. (31) and (32) as the alternative method for the parametrization. We confirm that the two methods give consistent results both for c_{sw} and K_c , and hence conclude that the systematic error due to the parameterization method for M and ΔM is small.

We fit c_{sw} to a rational function of g_0^2 and obtain the following interpolation formula

$$c_{\text{sw}} = \frac{1 - 0.194785g_0^2 - 0.110781g_0^4 - 0.0230239g_0^6 + 0.137401g_0^8}{1 - 0.460685g_0^2}. \quad (34)$$

The interpolation formula for K_c is obtained in a polynomial form

$$K_c = 1/8 + k^{(1)}g_0^2 + 0.000964911g_0^4 + 0.00298136g_0^6 + 0.00100995g_0^8 - 0.00235564g_0^{10}, \quad (35)$$

with $k^{(1)}=0.00843986$. These fits reproduce our data reasonably well with χ^2/dof of around 1.6. We also note that the coefficients of $O(g_0^2)$ terms in these fits are constrained so that these expressions coincide with their one-loop estimates [22, 23] up to $O(g_0^2)$.

We plot the β dependence of c_{SW} and K_c in Fig. 7. While c_{SW} in three-flavor QCD is well approximated by the one-loop estimate in the weak coupling region of $g_0^2 < 0.4$ ($\beta > 15$), it develops a significant deviation toward the strong coupling. Similar deviation is also observed in K_c . It is possible that these deviations are partly compensated by a better choice of the expansion parameter for lattice perturbation theory [24]. However, the large deviation at the strong coupling $g_0^2 \lesssim 1.0$ suggests that one-loop $O(a)$ -improved Wilson quark action leads to a significant $O(a)$ scaling violation in physical observables at the strong coupling region, where high statistics simulations are feasible with currently available computer resources. Therefore, the use of c_{SW} in Eq. (34) is essential to remove the $O(a)$ effects in practical lattice calculations.

C. N_f dependence

In Fig. 7, a comparison of our interpolation formula and those by the ALPHA Collaboration in two-flavor and quenched QCD suggests that c_{SW} monotonously decreases as N_f increases at fixed β . However, the difference between two and three-flavor QCD is not large. This comparison may also suffer from the systematic error due to the difference in the analysis method between the two collaborations. In order to study the N_f dependence more carefully, we determine c_{SW} and K_c for $N_f = 4, 2$ and 0 at $\beta = 9.6$ with an analysis method similar to that for $N_f = 3$. We obtain M and ΔM summarized in Tables XII–XIV. Fit to Eqs. (27) and (28) results in parameters given in Table XV and XVI. Using the improvement condition Eq. (25), we obtain c_{SW} and K_c summarized in Table XVII.

Figure 8 shows c_{SW} and K_c at $\beta = 9.6$ as a function of N_f . While K_c has an evident N_f dependence, that for c_{SW} is not so clear. The leading N_f dependence of c_{SW} is of order $g_0^4 N_f$. By fitting our results to a linear form in N_f

$$c_{\text{SW}} = c_0 + c_1 g_0^4 N_f, \quad (36)$$

we obtain $c_1 = -0.0117(40)$ which suggests that the two-loop N_f dependence is significant also in c_{SW} at the relatively weak coupling $\beta = 9.6$. If two-loop perturbative calculation of c_{SW} becomes available in future, it is interesting to compare the above estimate of c_1 to the perturbative estimate.

D. $O(a/L)$ uncertainty in c_{SW}

Our non-perturbative estimate of c_{SW} has $O(a/L)$ uncertainties, which affects physical observables at $O(a^2)$ when the spatial lattice size L is fixed. However, since we calculate c_{SW} with the spatial size in lattice units L/a fixed to a constant value 8, the $O(a/L)$ dependence of c_{SW} induces $O(a)$ effects in observables. These effects can be removed by extrapolating c_{SW} to the infinite volume limit, or interpolating to a fixed physical size in the whole region of g_0 . However, we are not able to do that in the present work, since our data are taken at a single lattice size at each g_0 .

We estimate the magnitude of $O(a/L)$ uncertainty in c_{SW} by using a modified improvement condition. In our study as well as in ALPHA's, the improvement condition Eq. (25) is adopted instead of $\Delta M = 0$ in order to remove the tree-level $O(a/L)$ correction to c_{SW} [5]. We extend this procedure to one-loop level, namely, the one-loop correction to ΔM for the finite lattice volume $8^3 \times 16$ given by [25]

$$a\Delta M^{(1)} = -(0.00004839 + 0.00006455 N_f) \quad (37)$$

is incorporated to the improvement condition

$$\begin{cases} M &= 0, \\ \Delta M &= \Delta M^{(0)} + g_0^2 \Delta M^{(1)} \end{cases} \quad (38)$$

in order to remove the $O(g_0^2 a/L)$ correction from c_{SW} .

From the parametrization of Eqs. (27)–(28) and the modified improvement condition Eq. (38), we obtain c_{SW} and K_c summarized in Table XVIII. A comparison with Table XI shows that the modified and original

improvement conditions give consistent results for c_{SW} with each other, and hence that the $O(g_0^2 a/L)$ correction is small in our results.

In Fig. 3, we observe that M and M' have a mild x_0 dependence at $4a < x_0 < 14a$. This may suggest that different choices of x_0 for the improvement conditions Eqs. (25) and (38) lead to a small difference in c_{SW} and hence its $O(a/L)$ ambiguity is not large.

From these observations, we expect that $O(g_0^2 a/L)$ corrections are not large at $L/a=8$, and that c_{SW} in the infinite volume limit is well approximated by our results. It is an important subject in future studies to confirm this point by a direct calculation of c_{SW} with varying L .

V. CONCLUSION

In this paper, we have performed a non-perturbative $O(a)$ -improvement of the Wilson quark action in three-flavor QCD with the plaquette gauge action. Our high statistics at carefully chosen simulation parameters c_{SW} and K enable us to determine non-perturbative c_{SW} with an accuracy of $\sim 5\%$ level in the wide range of $\beta = 12.0 - 5.2$. The main result of this study is the interpolation formula Eq. (34), with which $O(a)$ scaling violation in physical observables can be removed in future simulations at $\beta \geq 5.2$. As a byproduct, we also obtain the interpolation formula for K_c , which is useful to locate simulation points.

While it is expected that $O(a/L)$ uncertainty in c_{SW} is not large, this point should be confirmed by a direct determination of c_{SW} with varying L . A test of scaling properties of physical observables with our estimate of c_{SW} is an important next step toward an extensive simulation of three-flavor QCD with the plaquette gauge action.

We note that the range of β explored in this study is significantly higher than $\beta = 5.0$, where an artificial phase transition exists. Numerical studies at β below our range would possibly suffer from large distortion of scaling properties of physical observables. The use of improved gauge actions removes the lattice artifact and can push simulations toward coarser lattice spacings. This possibility is explored in a separate publication [9].

Acknowledgments

This work is supported by the Supercomputer Project No.98 (FY2003) of High Energy Accelerator Research Organization (KEK), and also in part by the Grant-in-Aid of the Ministry of Education (Nos. 12740133, 13135204, 13640260, 14046202, 14740173, 15204015, 15540251, 15540279, 16028201, 16540228, 16740147).

-
- [1] S. Duane, A.D. Kennedy, B.J. Pendleton, and D. Roweth, Phys. Lett. B **195**, 216 (1987); S. Gottlieb, W. Liu, D. Toussaint, R.L. Renken and R.L. Sugar, Phys. Rev. D **35**, 2531 (1987).
 - [2] M. Lüscher, Nucl. Phys. B **418**, 637 (1994).
 - [3] T. Takaishi and Ph. de Forcrand, hep-lat/0009024; Nucl. Phys. B (Proc.Suppl.) **94**, 818 (2001).
 - [4] B. Sheikholeslami and R. Wohlert, Nucl. Phys. B **259**, 572 (1985).
 - [5] M. Lüscher, S. Sint, R. Sommer, P. Weisz and U. Wolff, Nucl. Phys. B **491**, 323 (1997).
 - [6] K. Jansen and R. Sommer (ALPHA Collaboration), Nucl. Phys. B **530**, 185 (1998).
 - [7] S. Aoki *et al.* (JLQCD Collaboration), Nucl. Phys. B (Proc. Suppl.) **106**, 263 (2002).
 - [8] JLQCD Collaboration, in preparation.
 - [9] CP-PACS and JLQCD collaborations, in preparation.
 - [10] M. Lüscher, S. Sint, R. Sommer and P. Weisz, Nucl. Phys. B **478**, 365 (1996).
 - [11] C.R. Allton *et al.* (UKQCD Collaboration), Phys. Rev. D **65**, 054502 (2002).
 - [12] S. Aoki *et al.* (JLQCD Collaboration), Phys. Rev. D **68**, 054502 (2003).
 - [13] M. Lüscher, R. Narayanan, P. Weisz and U. Wolff, Nucl. Phys. B **384**, 168 (1992).
 - [14] M. Lüscher, R. Sommer, P. Weisz and U. Wolff, Nucl. Phys. B **389**, 247 (1993); S. Sint and R. Sommer, Nucl. Phys. B **465**, 71 (1996).
 - [15] X.-Q. Luo, Comput. Phys. Commun. **94**, 119 (1996).
 - [16] K. Jansen and C. Liu, Comput. Phys. Commun. **99**, 221 (1997).
 - [17] S. Aoki *et al.* (JLQCD Collaboration), Phys. Rev. D **65**, 094507 (2002).
 - [18] H. van der Vorst, SIAM J. Sc. Stat. Comp. **13**, 631 (1992).

- [19] R. Sommer *et al.* (ALPHA, CP-PACS and JLQCD Collaborations), Nucl. Phys. B (Proc. Suppl.) **129**, 405 (2004).
- [20] R. Sommer, Nucl. Phys. B **411**, 839 (1994).
- [21] A. D. Kennedy and J. Kuti, Phys. Rev. Lett. **54**, 2473 (1985).
- [22] R. Wohlert, DESY preprint 87-069 (1987), unpublished.
- [23] M. Lüscher and P. Weisz, Nucl. Phys. B **479**, 429 (1996).
- [24] G.P. Lepage and P.B. Mackenzie, Phys. Rev. D **48**, 2250 (1993).
- [25] S. Aoki, unpublished note.

TABLE I: Simulation parameters in two-flavor QCD at $\beta=5.2$.

c_{SW}	K	N_{traj}	c_{SW}	K	N_{traj}	c_{SW}	K	N_{traj}
1.50	0.14400	6400	1.90	0.13800	5600	2.02	0.13700	4800
1.50	0.14600	6200	1.90	0.13900	6387	2.30	0.13000	3200
1.80	0.13800	6400	2.02	0.13300	8000	2.30	0.13200	4000
1.80	0.14025	6400	2.02	0.13500	2176	3.00	0.12100	4000
1.90	0.13700	6400	2.02	0.13607	6400	3.00	0.12200	4000

TABLE II: Quark mass M and its difference ΔM in two-flavor QCD at $\beta=5.2$.

c_{SW}	K	aM	$a\Delta M$	c_{SW}	K	aM	$a\Delta M$	c_{SW}	K	aM	$a\Delta M$
1.50	0.14400	0.159(29)	-0.0039(44)	1.90	0.13800	0.0083(22)	-0.0008(16)	2.02	0.13700	-0.0411(17)	0.0040(27)
1.50	0.14600	-0.0012(39)	0.0073(24)	1.90	0.13900	-0.0338(23)	0.0033(14)	2.30	0.13000	0.0546(10)	-0.00108(91)
1.80	0.13800	0.136(10)	-0.0073(28)	2.02	0.13300	0.1361(51)	-0.0003(12)	2.30	0.13200	-0.0187(11)	-0.00296(97)
1.80	0.14025	-0.0048(29)	0.0024(22)	2.02	0.13500	0.0541(47)	0.0002(15)	3.00	0.12100	0.03767(84)	-0.00782(49)
1.90	0.13700	0.0629(31)	-0.0038(16)	2.02	0.13607	0.0005(20)	-0.0022(14)	3.00	0.12200	0.00455(71)	-0.00735(82)

TABLE III: Fit parameters for Eqs. (27) and (28) in two-flavor QCD at $\beta=5.2$.

χ^2/dof	a_M	$b_M^{(1)}$	$b_M^{(2)}$	$c_M^{(1)}$	$c_M^{(2)}$	d_M
6.37	-28.8(7.1)	8.9(2.5)	-0.67(22)	-6.3(2.2)	-0.15(17)	0.83(39)
χ^2/dof	$a_{\Delta M}$	$b_{\Delta M}^{(1)}$	$b_{\Delta M}^{(2)}$	$c_{\Delta M}^{(1)}$	$c_{\Delta M}^{(2)}$	$d_{\Delta M}$
1.88	11.2(3.2)	-3.9(1.1)	0.336(98)	3.16(95)	0.214(71)	-0.54(17)

TABLE IV: Simulation parameters in three-flavor QCD.

$\beta = 12.0$							
$c_{\text{SW}} = 1.08624$		$c_{\text{SW}} = 1.12036$		$c_{\text{SW}} = 1.15448$		$c_{\text{SW}} = 1.18860$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.128958	3000	0.128958	3000	0.128958	3000	0.128958	3000
0.129607	3000	0.129607	3000	0.129607	3000	0.129607	3000
0.130257	3000	0.130257	3000	0.130257	3000	0.130257	3000
0.130906	3000	0.130906	3000	0.130906	3000	0.130906	3000
$\beta = 9.6$							
$c_{\text{SW}} = 1.12990$		$c_{\text{SW}} = 1.16539$		$c_{\text{SW}} = 1.20089$		$c_{\text{SW}} = 1.23638$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13043	7000	0.13043	7000	0.13043	7000	0.13043	7000
0.13109	7000	0.13109	7000	0.13109	7000	0.13109	7000
0.13175	7000	0.13175	7000	0.13175	7000	0.13175	7000
0.13240	7000	0.13240	7000	0.13230	1800	0.13230	7000
$\beta = 7.4$							
$c_{\text{SW}} = 1.2258$		$c_{\text{SW}} = 1.2643$		$c_{\text{SW}} = 1.3028$		$c_{\text{SW}} = 1.3413$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13293	5400	0.13293	5400	0.13293	5400	0.13293	5400
0.13360	5400	0.13360	5400	0.13360	5400	0.13360	5400
0.13427	5400	0.13427	5400	0.13427	5400	0.13393	400
0.13494	5400	0.13470	3400	–	–	0.13427	5400
$\beta = 6.8$							
$c_{\text{SW}} = 1.2783$		$c_{\text{SW}} = 1.3184$		$c_{\text{SW}} = 1.3586$		$c_{\text{SW}} = 1.3987$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13391	4200	0.13391	4200	0.13391	4200	0.13391	4200
0.13459	4200	0.13459	4200	0.13459	4200	0.13459	4200
0.13526	4200	0.13526	4200	0.13526	4200	0.13492	3700
0.13594	4200	0.13560	4200	0.13540	3700	0.13500	3150
$\beta = 6.3$							
$c_{\text{SW}} = 1.3117$		$c_{\text{SW}} = 1.3675$		$c_{\text{SW}} = 1.4233$		$c_{\text{SW}} = 1.4791$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13501	4600	0.13501	4600	0.13446	4600	0.13446	4600
0.13555	4600	0.13555	4600	0.13501	4600	0.13501	4600
0.13609	4600	0.13609	4600	0.13555	4600	0.13530	1100
0.13664	4600	0.13664	4600	0.13609	4600	0.13555	4600
$\beta = 6.0$							
$c_{\text{SW}} = 1.3237$		$c_{\text{SW}} = 1.3801$		$c_{\text{SW}} = 1.4364$		$c_{\text{SW}} = 1.4927$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13552	4200	0.13552	4200	0.13498	4200	0.13498	4200
0.13607	4200	0.13607	4200	0.13552	4200	0.13552	4200
0.13661	4200	0.13661	4200	0.13607	4200	0.13607	4200
0.13716	4200	0.13716	4200	0.13661	4200	0.13634	4200
$\beta = 5.7$							
$c_{\text{SW}} = 1.5431$		$c_{\text{SW}} = 1.6088$		$c_{\text{SW}} = 1.6745$		$c_{\text{SW}} = 1.7401$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13575	5000	0.13541	5000	0.13507	5000	0.13400	4500
0.13644	5000	0.13610	5000	0.13540	4400	0.13450	4100
0.13712	4700	0.13644	3700	0.13575	5000	0.13473	5000
0.13750	4500	0.13678	5000	0.13585	2700	0.13500	4400

TABLE IV: (*Continued*)

$\beta=5.4$							
$c_{\text{SW}}=1.6$		$c_{\text{SW}}=1.7$		$c_{\text{SW}}=1.8$		$c_{\text{SW}}=1.9$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13750	9000	0.13600	9000	0.13480	9000	0.13330	9000
0.13810	9000	0.13630	7900	0.13520	9000	0.13370	9000
0.13825	7100	0.13660	9000	0.13560	9000	0.13410	9000
0.13840	8200	0.13720	9000	0.13580	7700	0.13450	9000

$\beta=5.2$							
$c_{\text{SW}}=1.70$		$c_{\text{SW}}=1.85$		$c_{\text{SW}}=2.00$		$c_{\text{SW}}=2.15$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.1375	7800	0.1357	7800	0.1336	7800	0.1316	7800
0.1379	7800	0.1361	7800	0.1340	7800	0.1320	7800
0.1383	7800	0.1365	7800	0.1344	7800	0.1322	5500
0.1387	6300	0.1366	7400	0.1345	7400	0.1324	7800

TABLE V: Simulation parameters in four-flavor QCD at $\beta=9.6$.

$c_{\text{SW}}=1.12990$		$c_{\text{SW}}=1.16539$		$c_{\text{SW}}=1.20089$		$c_{\text{SW}}=1.23638$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13043	5000	0.13043	5000	0.13043	5000	0.13043	5000
0.13109	5000	0.13109	5000	0.13109	5000	0.13109	5000
0.13175	5000	0.13175	5000	0.13175	5000	0.13175	5000
0.13240	5000	0.13240	5000	0.13240	5000	0.13240	5000

TABLE VI: Simulation parameters in two-flavor QCD at $\beta=9.6$.

$c_{\text{SW}}=1.12990$		$c_{\text{SW}}=1.16539$		$c_{\text{SW}}=1.20089$		$c_{\text{SW}}=1.23638$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13043	5000	0.13043	5000	0.13043	5000	0.13043	5000
0.13109	5000	0.13109	5000	0.13109	5000	0.13109	5000
0.13175	5000	0.13175	5000	0.13175	5000	0.13175	5000
0.13240	5000	0.13240	5000	0.13240	5000	0.13240	5000

TABLE VII: Simulation parameters in quenched QCD at $\beta=9.6$.

$c_{\text{SW}}=1.12990$		$c_{\text{SW}}=1.16539$		$c_{\text{SW}}=1.20089$		$c_{\text{SW}}=1.23638$	
K	N_{traj}	K	N_{traj}	K	N_{traj}	K	N_{traj}
0.13043	8000	0.13043	8000	0.13043	8000	0.13043	8000
0.13109	8000	0.13109	8000	0.13109	8000	0.13109	8000
0.13175	8000	0.13175	8000	0.13175	8000	0.13175	8000
0.13240	8000	0.13240	8000	0.13240	8000	0.13240	8000

TABLE VIII: Quark mass M and its difference ΔM in three-flavor QCD.

$\beta = 12.0$					
$c_{\text{sw}} = 1.08624$			$c_{\text{sw}} = 1.12036$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.128958	0.03502(11)	0.00107(13)	0.128958	0.03066(8)	0.00067(10)
0.129607	0.01498(13)	0.00099(12)	0.129607	0.01028(10)	0.00072(16)
0.130257	-0.00544(10)	0.00100(14)	0.130257	-0.01011(12)	0.00040(12)
0.130906	-0.02591(11)	0.00116(12)	0.130906	-0.03088(11)	0.00052(12)
$c_{\text{sw}} = 1.15448$			$c_{\text{sw}} = 1.18860$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.128958	0.02581(9)	0.00027(12)	0.128958	0.02125(9)	-0.00029(11)
0.129607	0.00561(11)	0.00015(11)	0.129607	0.00076(10)	-0.00027(11)
0.130257	-0.01512(14)	0.00008(13)	0.130257	-0.01965(11)	-0.00044(13)
0.130906	-0.03555(17)	0.00025(13)	0.130906	-0.04041(12)	-0.00043(14)
$\beta = 9.6$					
$c_{\text{sw}} = 1.12990$			$c_{\text{sw}} = 1.16539$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13043	0.03818(12)	0.00118(10)	0.13043	0.03226(9)	0.00067(10)
0.13109	0.01805(9)	0.00100(12)	0.13109	0.01187(10)	0.00062(12)
0.13175	-0.00275(10)	0.00104(14)	0.13175	-0.00871(12)	0.00056(13)
0.13240	-0.02312(11)	0.00104(13)	0.13240	-0.02948(11)	0.00063(16)
$c_{\text{sw}} = 1.20089$			$c_{\text{sw}} = 1.23638$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13043	0.02596(11)	0.00037(11)	0.13043	0.01995(9)	-0.00037(14)
0.13109	0.00559(12)	0.00014(13)	0.13109	-0.00091(10)	-0.00012(9)
0.13175	-0.01522(11)	0.00005(12)	0.13175	-0.02145 (11)	-0.00044(13)
0.13230	-0.03256(19)	0.00033(24)	0.13230	-0.03934 (13)	-0.00054(14)
$\beta = 7.4$					
$c_{\text{sw}} = 1.2258$			$c_{\text{sw}} = 1.2643$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13293	0.04147(22)	0.00135(19)	0.13293	0.03265(15)	0.00067(19)
0.13360	0.02134(21)	0.00158(25)	0.13360	0.01209(23)	0.00065(25)
0.13427	0.00079(21)	0.00099(26)	0.13427	-0.00942(21)	0.00102(23)
0.13494	-0.02065(24)	0.00077(28)	0.13470	-0.02292(28)	0.00089(34)
$c_{\text{sw}} = 1.3028$			$c_{\text{sw}} = 1.3413$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13293	0.02296(25)	0.00035(24)	0.13293	0.01309(22)	0.00004(23)
0.13360	0.00214(17)	0.00018(19)	0.13360	-0.00756(21)	0.00029(28)
0.13427	-0.01888(23)	0.00052(22)	0.13393	-0.01774(64)	-0.00018(68)
-	-	-	0.13427	-0.02867(26)	-0.00048(36)
$\beta = 6.8$					
$c_{\text{sw}} = 1.2783$			$c_{\text{sw}} = 1.3184$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13391	0.04462(36)	0.00066(28)	0.13391	0.03342(33)	0.00092(28)
0.13459	0.02457(29)	0.00079(32)	0.13459	0.01358(28)	0.00051(27)
0.13526	0.00338(33)	0.00098(30)	0.13526	-0.00823(30)	0.00065(40)
0.13594	-0.01824(30)	0.00163(37)	0.13560	-0.01839(36)	0.00054(39)
$c_{\text{sw}} = 1.3586$			$c_{\text{sw}} = 1.3987$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13391	0.02240(31)	0.00049(26)	0.13391	0.01208(22)	-0.00021(35)
0.13459	0.00175(31)	0.00031(23)	0.13459	-0.00953(30)	-0.00030(27)
0.13526	-0.01924(27)	0.00033(34)	0.13492	-0.01956(58)	0.00025(45)
0.13540	-0.02474(30)	0.00058(46)	0.13500	-0.02271(43)	-0.00014(34)

TABLE VIII: (*Continued*)

$\beta=6.3$					
$c_{\text{SW}} = 1.3117$			$c_{\text{SW}} = 1.3675$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13501	0.05392(55)	0.00085(50)	0.13501	0.03677(44)	0.00097(29)
0.13555	0.03766(40)	0.00107(32)	0.13555	0.02065(30)	0.00063(33)
0.13609	0.02152(36)	0.00080(38)	0.13609	0.00353(36)	0.00058(32)
0.13664	0.00403(41)	0.00175(34)	0.13664	-0.01345(53)	0.00168(41)
$c_{\text{SW}} = 1.4233$			$c_{\text{SW}} = 1.4791$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13446	0.03635(37)	0.00070(37)	0.13446	0.01834(37)	-0.00034(33)
0.13501	0.01915(34)	0.00028(37)	0.13501	0.00198(38)	0.00006(30)
0.13555	0.00284(43)	0.00092(33)	0.13530	-0.00807(57)	0.00075(47)
0.13609	-0.01497(43)	0.00050(37)	0.13555	-0.01599(41)	-0.00039(51)
$\beta=6.0$					
$c_{\text{SW}} = 1.3237$			$c_{\text{SW}} = 1.3801$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13552	0.07608(51)	0.00154(39)	0.13552	0.05826(50)	0.00156(37)
0.13607	0.06056(57)	0.00175(43)	0.13607	0.04182(39)	0.00124(44)
0.13661	0.04411(52)	0.00120(48)	0.13661	0.02400(46)	0.00077(40)
0.13716	0.02755(53)	0.00241(53)	0.13716	0.00843(50)	0.00259(53)
$c_{\text{SW}} = 1.4364$			$c_{\text{SW}} = 1.4927$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13498	0.05492(48)	0.00054(40)	0.13498	0.03635(37)	-0.00060(60)
0.13552	0.03893(49)	0.00035(54)	0.13552	0.01878(41)	0.00030(45)
0.13607	0.02221(48)	0.00107(44)	0.13607	0.00235(45)	-0.00005(48)
0.13661	0.00562(49)	0.00031(50)	0.13634	-0.00692(49)	0.00076(39)
$\beta=5.7$					
$c_{\text{SW}} = 1.5431$			$c_{\text{SW}} = 1.6088$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13575	0.04354(59)	0.00079(42)	0.13541	0.02748(50)	-0.00068(47)
0.13644	0.02091(55)	-0.00039(51)	0.13610	0.00695(130)	0.00053(54)
0.13712	-0.00181(57)	0.00072(53)	0.13644	-0.00587(70)	-0.00087(64)
0.13750	-0.01386(70)	0.00007(72)	0.13678	-0.01702(60)	-0.00089(65)
$c_{\text{SW}} = 1.6745$			$c_{\text{SW}} = 1.7401$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13507	0.01331(69)	-0.00081(51)	0.13400	0.02229(42)	-0.00140(52)
0.13540	0.00278(64)	-0.00022(61)	0.13450	0.00521(59)	-0.00104(39)
0.13575	-0.00947(54)	-0.00078(43)	0.13473	-0.00173(44)	-0.00082(42)
0.13585	-0.01124(62)	-0.00115(51)	0.13500	-0.01153(59)	-0.00172(69)
$\beta=5.4$					
$c_{\text{SW}} = 1.6$			$c_{\text{SW}} = 1.7$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13750	0.03310(70)	0.00015(43)	0.13600	0.03592(56)	-0.00031(40)
0.13810	0.01245(75)	0.00083(52)	0.13630	0.02622(54)	0.00032(52)
0.13825	0.00684(78)	0.00053(62)	0.13660	0.01653(48)	0.00030(46)
0.13840	0.00382(124)	0.00100(49)	0.13720	-0.00589(52)	0.00106(57)
$c_{\text{SW}} = 1.8$			$c_{\text{SW}} = 1.9$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13480	0.03073(51)	-0.00129(56)	0.13330	0.03600(40)	-0.00149(40)
0.13520	0.01650(50)	-0.00048(49)	0.13370	0.02250(34)	-0.00167(51)
0.13560	0.00373(52)	0.00010(47)	0.13410	0.00871(47)	-0.00110(40)
0.13580	-0.00426(70)	-0.00062(53)	0.13450	-0.00508(51)	-0.00194(51)

TABLE VIII: (*Continued*)

$\beta = 5.2$					
$c_{\text{SW}} = 1.70$			$c_{\text{SW}} = 1.85$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.1375	0.05191(131)	0.00102(87)	0.1357	0.03406(87)	-0.00004(54)
0.1379	0.03776(89)	0.00062(67)	0.1361	0.01844(73)	0.00045(46)
0.1383	0.02311(82)	0.00196(54)	0.1365	0.00287(78)	0.00019(81)
0.1387	0.00807(123)	0.00040(110)	0.1366	0.00037(84)	-0.00038(56)
$c_{\text{SW}} = 2.00$			$c_{\text{SW}} = 2.15$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.1336	0.02982(64)	-0.00161(43)	0.1316	0.02568(70)	-0.00314(65)
0.1340	0.01467(74)	-0.00089(48)	0.1320	0.01350(49)	-0.00281(46)
0.1344	-0.00003(68)	-0.00219(48)	0.1322	0.00364(74)	-0.00348(77)
0.1345	-0.00405(69)	-0.00133(48)	0.1324	-0.00202(51)	-0.00284(60)

TABLE IX: Fit parameters for Eq. (27) in three-flavor QCD.

β	χ^2/dof	a_M	$b_M^{(1)}$	$b_M^{(2)}$	$c_M^{(1)}$	$c_M^{(2)}$	d_M
12.0	0.96	-10.9(1.1)	2.40(28)	-0.125(18)	-0.47(13)	-0.012(23)	0.046(16)
9.6	1.41	-13.1(1.1)	3.02(30)	-0.168(20)	-0.64(14)	-0.047(22)	0.076(17)
7.4	1.17	-15.9(2.9)	3.90(81)	-0.233(55)	-0.90(34)	-0.057(39)	0.106(46)
6.8	1.57	-24.9(4.3)	6.6(1.2)	-0.430(85)	-2.30(47)	0.016(53)	0.266(68)
6.3	0.66	-19.0(5.8)	5.0(1.7)	-0.33(12)	-1.99(66)	-0.094(40)	0.262(97)
6.0	1.19	-12.2(6.6)	3.2(1.9)	-0.20(13)	-1.75(64)	-0.111(40)	0.234(92)
5.7	1.16	-7.6(9.5)	1.6(3.0)	-0.07(23)	-0.4(1.8)	-0.007(99)	0.005(280)
5.4	0.92	-27(11)	8.0(3.7)	-0.58(30)	-5.0(2.7)	-0.19(17)	0.70(45)
5.2	1.42	-38(22)	11.9(7.4)	-0.91(62)	-8.0(5.5)	-0.32(35)	1.17(93)

TABLE X: Fit parameters for Eq. (28) in three-flavor QCD.

β	χ^2/dof	$a_{\Delta M}$	$b_{\Delta M}^{(1)}$	$b_{\Delta M}^{(2)}$	$c_{\Delta M}^{(1)}$	$c_{\Delta M}^{(2)}$	$d_{\Delta M}$
12.0	0.59	1.6(1.2)	-0.41(32)	0.025(21)	-0.12(16)	-0.008(26)	0.016(19)
9.6	0.91	0.6(1.3)	-0.14(35)	0.008(23)	-0.14(16)	-0.007(25)	0.019(19)
7.4	1.13	-3.1(3.4)	0.82(93)	-0.053(64)	0.09(38)	0.025(44)	-0.022(51)
6.8	0.40	6.2(4.4)	-1.7(1.2)	0.109(84)	-0.06(50)	-0.051(49)	0.025(66)
6.3	1.14	7.1(5.3)	-2.0(1.5)	0.14(11)	0.49(61)	-0.012(38)	-0.062(91)
6.0	1.28	8.7(6.3)	-2.5(1.8)	0.18(13)	0.78(60)	0.003(37)	-0.108(84)
5.7	1.06	-4.8(8.7)	1.5(2.7)	-0.11(21)	-0.7(1.5)	0.008(80)	0.09(23)
5.4	0.53	-6.9(10.8)	2.4(3.6)	-0.21(29)	-2.1(2.6)	-0.18(16)	0.38(43)
5.2	0.67	-26(16)	8.7(5.4)	-0.73(45)	-6.6(4.0)	-0.42(25)	1.10(67)

TABLE XI: Non-perturbative estimate of c_{SW} and K_c in three-flavor QCD obtained from tree-level improvement condition Eq. (25).

β	c_{SW}	K_c
12.0	1.1415(48)	0.129841(21)
9.6	1.1916(50)	0.131321(28)
7.4	1.316(11)	0.133567(87)
6.8	1.358(13)	0.13466(12)
6.3	1.447(15)	0.13539(16)
6.0	1.494(14)	0.13612(16)
5.7	1.544(32)	0.13706(39)
5.4	1.740(30)	0.13650(41)
5.2	1.764(103)	0.13789(159)

TABLE XII: Quark mass M and its difference ΔM in four-flavor QCD at $\beta=9.6$.

$c_{\text{SW}} = 1.12990$			$c_{\text{SW}} = 1.16539$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13043	0.03530(11)	0.00111(14)	0.13043	0.02955(11)	0.00050(11)
0.13109	0.01527(12)	0.00118(15)	0.13109	0.00892(12)	0.00068(13)
0.13175	-0.00522(12)	0.00110(12)	0.13175	-0.01158(11)	0.00069(12)
0.13240	-0.02611(13)	0.00123(12)	0.13240	-0.03256(14)	0.00061(13)
$c_{\text{SW}} = 1.20089$			$c_{\text{SW}} = 1.23638$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13043	0.02328(12)	0.00018(11)	0.13043	0.01723(12)	-0.00032(14)
0.13109	0.00280(14)	0.00028(13)	0.13109	-0.00364(12)	-0.00021(11)
0.13175	-0.01825(10)	0.00028(15)	0.13175	-0.02462(17)	-0.00035(15)
0.13240	-0.03903(15)	0.00004(15)	0.13240	-0.04592(14)	-0.00023(13)

TABLE XIII: Quark mass M and its difference ΔM in two-flavor QCD at $\beta=9.6$.

$c_{\text{SW}} = 1.12990$			$c_{\text{SW}} = 1.16539$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13043	0.04089(10)	0.00127(13)	0.13043	0.03508(10)	0.00085(12)
0.13109	0.02085(10)	0.00119(13)	0.13109	0.01482(12)	0.00053(16)
0.13175	0.00047(12)	0.00102(16)	0.13175	-0.00585(13)	0.00070(14)
0.13240	-0.01998(13)	0.00104(13)	0.13240	-0.02646(15)	0.00068(15)
$c_{\text{SW}} = 1.20089$			$c_{\text{SW}} = 1.23638$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13043	0.02884(12)	0.00013(13)	0.13043	0.02259(10)	-0.00021(13)
0.13109	0.00855(11)	0.00045(14)	0.13109	0.00218(10)	-0.00003(12)
0.13175	-0.01225(13)	0.00013(19)	0.13175	-0.01857(13)	-0.00000(13)
0.13240	-0.03270(12)	0.00032(22)	0.13240	-0.03927(11)	-0.00024(16)

TABLE XIV: Quark mass M and its difference ΔM in quenched QCD at $\beta=9.6$.

$c_{\text{SW}} = 1.12990$			$c_{\text{SW}} = 1.16539$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13043	0.04646(7)	0.00141(10)	0.13043	0.04059(8)	0.00095(11)
0.13109	0.02679(8)	0.00131(11)	0.13109	0.02064(8)	0.00080(11)
0.13175	0.00656(7)	0.00131(14)	0.13175	0.00041(7)	0.00084(10)
0.13240	-0.01324(8)	0.00132(10)	0.13240	-0.01978(8)	0.00074(11)
$c_{\text{SW}} = 1.20089$			$c_{\text{SW}} = 1.23638$		
K	aM	$a\Delta M$	K	aM	$a\Delta M$
0.13043	0.03458(9)	0.00036(11)	0.13043	0.02840(8)	-0.00015(10)
0.13109	0.01451(7)	0.00034(12)	0.13109	0.00818(7)	0.00006(10)
0.13175	-0.00580(8)	0.00050(14)	0.13175	-0.01231(8)	-0.00005(13)
0.13240	-0.02607(10)	0.00053(16)	0.13240	-0.03247(10)	-0.00013(21)

TABLE XV: Fit parameters in Eq. (27) for $N_f=4, 2$ and 0.

N_f	β	χ^2/dof	a_M	$b_M^{(1)}$	$b_M^{(2)}$	$c_M^{(1)}$	$c_M^{(2)}$	d_M
4	9.6	0.96	-14.7(1.3)	3.52(33)	-0.206(22)	-1.14(15)	-0.041(25)	0.140(19)
2	9.6	0.48	-13.5(1.2)	3.15(31)	-0.177(20)	-0.64(14)	-0.052(23)	0.077(17)
0	9.6	1.03	-11.71(80)	2.70(21)	-0.150(14)	-0.80(10)	-0.040(16)	0.094(12)

TABLE XVI: Fit parameters in Eq. (28) for $N_f=4, 2$ and 0.

N_f	β	χ^2/dof	$a_{\Delta M}$	$b_{\Delta M}^{(1)}$	$b_{\Delta M}^{(2)}$	$c_{\Delta M}^{(1)}$	$c_{\Delta M}^{(2)}$	$d_{\Delta M}$
4	9.6	0.42	-1.2(1.3)	0.33(34)	-0.023(22)	-0.14(16)	0.015(26)	0.012(19)
2	9.6	0.80	-1.0(1.4)	0.24(38)	-0.014(25)	0.15(17)	0.013(28)	-0.026(21)
0	9.6	0.53	-0.3(1.2)	0.06(31)	-0.002(20)	0.11(14)	0.012(23)	-0.020(18)

TABLE XVII: Non-perturbative estimate of c_{sw} and K_c for $N_f = 4, 2$ and 0 obtained from tree-level improvement condition Eq. (25).

N_f	β	c_{sw}	K_c
4	9.6	1.1954(48)	0.131210(27)
2	9.6	1.2028(63)	0.131353(36)
0	9.6	1.2112(44)	0.131502(25)

TABLE XVIII: Non-perturbative estimate of c_{sw} and K_c in three-flavor QCD obtained from one-loop level improvement condition Eq. (38).

β	c_{sw}	K_c
12.0	1.1507(46)	0.129801(20)
9.6	1.2039(46)	0.131252(26)
7.4	1.336(14)	0.13340(12)
6.8	1.378(12)	0.13447(11)
6.3	1.470(16)	0.13515(17)
6.0	1.508(17)	0.13596(20)
5.7	1.569(27)	0.13676(32)
5.4	1.770(26)	0.13609(35)
5.2	1.811(67)	0.1372(10)




FIG. 1: Plots of M (top figure) and ΔM (bottom figure) in two-flavor QCD at $\beta = 5.2$ as a function of $1/K$. The bottom figure shows ΔM as a function of M .




FIG. 2: Mass difference ΔM at $M=0$ as a function of c_{sw} . Filled and open symbols are obtained from the fit Eq. (31) and combined fit Eqs. (27) and (28). Results from the combined fit are slightly shifted in the horizontal direction for a better visibility. Dotted and dashed lines show linear and quadratic fit Eq. (32). Dot-dashed line shows the tree-level value $\Delta M^{(0)} = 0.000277$.




FIG. 3: Quark masses M and M' in three-flavor QCD as a function of x_0 . Top, middle and bottom figures show data at $\beta=12.0$, 9.6 and 5.2 , respectively.




FIG. 4: Quark mass M as a function of $1/K$. Top, middle and bottom figures show data at $\beta = 12.0$, 9.6 and 5.2 , respectively. Dotted lines are fit curve of Eq. (27).




FIG. 5: Mass difference ΔM as a function of $1/K$. Top, middle and bottom figures show data at $\beta=12.0$, 9.6 and 5.2, respectively. Dotted lines are fit curve of Eq. (28).




FIG. 6: Mass difference ΔM as a function of M in three-flavor QCD. Top, middle and bottom figures show data at $\beta=12.0, 9.6$ and 5.2 , respectively. Dotted lines are fit curve reproduced from Eqs. (27) and (28).




FIG. 7: Non-perturbatively determined c_{SW} (top figure) and K_c (bottom figure) as a function of g_0^2 . For K_c in quenched QCD, author's interpolation of the ALPHA's results is plotted.




FIG. 8: Plots of c_{SW} (top figure), K_c (bottom figure) at $\beta=9.6$ as a function of N_f . Dashed line shows linear fit to data.